

## **Safety of Medicines and the use of animals in research**

We are concerned about the misleading claims made by Kathy Archibald and colleagues in their Correspondence (June 4, p 1915).

The main claim made is that more medicines are failing, that adverse drug reactions have increased, and that these problems are caused by inadequacies in the medicines development process. They blamed an “over-reliance of the pharmaceutical industry on the use of animals to predict drug behaviour in man”<sup>1</sup>.

It is essential for all doctors, patients, regulators and the bioscience sector globally that new medicines are safe and effective. Sound and sophisticated science is used at all stages of research into health and disease, and in the development of new treatments. Animal research and testing is a small but vital part of this effort, and does not represent, in any sense, “over-reliance”.

Animal research and testing is essential for understanding normal and disease processes. In the preclinical development of new treatment candidates, animals are essential for understanding their pharmacokinetics and pharmacodynamics, and for detecting unforeseen toxic effects. The identification of developmental and carcinogenic hazards is particularly reliant on continued animal testing, since such effects would not become evident in patients for many years.

Nobody is permitted to use animals where there is a viable alternative<sup>2</sup>. All parties support the significant investment in the search for alternatives to animals in research and ways of improving their welfare and reducing the numbers used. The UK leads in this area<sup>3</sup>. The global pharmaceutical industry is also making strides to encourage harmonisation of international regulatory requirements to minimise animal testing wherever possible<sup>4</sup>.

The UK’s Medicines and Healthcare Products Regulatory Agency has acknowledged that non-animal testing is used wherever possible, but adds “at present there are no laboratory methods available to totally replace animal testing of medicines” (personal communication).

99% of drug candidates discovered at the beginning of the process are indeed eliminated, most during early research phases, but this is a necessary part of the process of demonstrating efficacy as well as safety<sup>6</sup>. More relevant is the success of the vast majority of medicines approved for market by the regulators. Only a very small number – estimated at less than 1% – are subsequently withdrawn owing to serious side-effects.

93% percent of people admitted to hospital with an adverse drug reaction have a Type A response. This means that the side-effects are predictable and could have been avoided if the medicine had been taken according to its prescribing information, or if the prescriber had been aware of other medicines being taken by the patient<sup>7</sup>. Most adverse drug reactions are, therefore, avoidable rather than the result of inadequacies in the drug development process.

Millions of people are alive today thanks to medical advances. Despite these efforts there are still diseases without adequate treatments – for example dementia, many cancers and heart failure. We all need to strive to improve every step of the development process, but this does not mean we should tear up a tried and tested process that has helped so many patients. Very recently, a research on mice, stem cells and a new polypeptide drug showed how the heart may be able to repair itself<sup>8</sup>. It is difficult to see how this advance could have been achieved without animal research.

Frances Balkwill, Barts Cancer Institute, Queen Mary University of London and Understanding Animal Research

Stephen Whitehead, Association of the British Pharmaceutical Industry

Phil Willis Lord Willis of Knaresborough, Association of Medical Research Charities

Nigel Gaymond, BioIndustry Association

Alastair Kent, Genetic Alliance UK

Clive Page, Animal Science Group, Society of Biology

Robin Lovell-Badge, Institute of Animal Technology

Roger Morris, School of Biomedical Sciences, King's College London

Roger Lemon, UCL Institute of Neurology

Duncan Banks, British Neuroscience Association

Corresponding author: Frances Balkwill, f.balkwill@qmul.ac.uk

1. Archibald K, Coleman R, Foster C, on behalf of 19 other signatories. Open letter to UK Prime Minister David Cameron and Health Secretary Andrew Lansley on safety of medicines. *Lancet* 2011; **377**: 1915.
2. House of Commons. Guidance on the Operation of the Animals (Scientific Procedures) Act 1986. London: Stationery Office, 2000.
3. National Centre for the Replacement, Refinement and Reduction of Animals in Research, Annual Report 2010. London: NC3Rs 2010.  
(<http://www.nc3rs.org.uk/document.asp?id=1456>) (accessed June 24, 2011).
4. International Conference on Harmonisation of Technical Requirements for Registration of Pharmaceuticals for Human Use. Vision: about ICH <http://www.ich.org/about/vision.html> (accessed June 9, 2011).
5. Paul S, Mytelka DS, Dunwiddie CT et al. How to improve R&D productivity; the pharmaceutical industry's grand challenge. *Nat Rev Drug Discov* 2010; **9** 203–14.
6. Pirmohamed M, James S, Meakin S, et al (2004). Adverse drug reactions as cause of admission to hospital: prospective analysis of 18,820 patients. *BMJ* 2004; **329**, 5–19.
7. Smart N, Bollini S, Dubé KN, et al. De novo cardiomyocytes from within the activated adult heart after injury. *Nature* 2011 doi:10.1038/nature10188.